

# CM-Points on Straight Lines

A joint work with **Amalia Pizarro-Madariaga**

Bill Allombert & Yuri Bilu

Bordeaux

September 23, 2014

## Complex Multiplication

Lattices

$j$ -invariant

Complex Multiplication

Class Field Theory

The Class Number

## Theorem of André

Special Points and Special Curves

Theorem of André

## CM-Points on Straight Lines

Kühne's "uniformity observation"

CM-Points on Straight Lines

## The Proof

Equality of CM-fields

The Proof

## Proof of Theorem ECMF

Discriminants with Class Group Annihilated by 2

Proof of Theorem ECMF

## Complex Multiplication

Lattices

$j$ -invariant

Complex Multiplication

Class Field Theory

The Class Number

## Theorem of André

Special Points and Special Curves

Theorem of André

## CM-Points on Straight Lines

Kühne's "uniformity observation"

CM-Points on Straight Lines

## The Proof

Equality of CM-fields

The Proof

## Proof of Theorem ECMF

Discriminants with Class Group Annihilated by 2

Proof of Theorem ECMF

# Lattices

**Lattice** in  $\mathbb{C}$ : a discrete (additive) group of rank 2

# Lattices

**Lattice** in  $\mathbb{C}$ : a discrete (additive) group of rank 2

**Example:**  $\{a + bi : a, b \in \mathbb{Z}\} = \langle i, 1 \rangle$ .

# Lattices

**Lattice** in  $\mathbb{C}$ : a discrete (additive) group of rank 2

**Example:**  $\{a + bi : a, b \in \mathbb{Z}\} = \langle i, 1 \rangle$ .

**Isomorphism of lattices:**  $\Lambda \cong \Lambda'$  if  $\Lambda' = \alpha\Lambda$ ,  $\alpha \in \mathbb{C}$ .

# Lattices

**Lattice** in  $\mathbb{C}$ : a discrete (additive) group of rank 2

**Example:**  $\{a + bi : a, b \in \mathbb{Z}\} = \langle i, 1 \rangle$ .

**Isomorphism of lattices:**  $\Lambda \cong \Lambda'$  if  $\Lambda' = \alpha\Lambda$ ,  $\alpha \in \mathbb{C}$ .

- ▶ Every lattice is isomorphic to  $\langle \tau, 1 \rangle$ ,  $\text{Im } \tau > 0$

# Lattices

**Lattice** in  $\mathbb{C}$ : a discrete (additive) group of rank 2

**Example:**  $\{a + bi : a, b \in \mathbb{Z}\} = \langle i, 1 \rangle$ .

**Isomorphism of lattices:**  $\Lambda \cong \Lambda'$  if  $\Lambda' = \alpha\Lambda$ ,  $\alpha \in \mathbb{C}$ .

- ▶ Every lattice is isomorphic to  $\langle \tau, 1 \rangle$ ,  $\text{Im } \tau > 0$



# Lattices

**Lattice** in  $\mathbb{C}$ : a discrete (additive) group of rank 2

**Example:**  $\{a + bi : a, b \in \mathbb{Z}\} = \langle i, 1 \rangle$ .

**Isomorphism of lattices:**  $\Lambda \cong \Lambda'$  if  $\Lambda' = \alpha\Lambda$ ,  $\alpha \in \mathbb{C}$ .

- ▶ Every lattice is isomorphic to  $\langle \tau, 1 \rangle$ ,  $\text{Im } \tau > 0$
- ▶  $\langle \tau, 1 \rangle \cong \langle \tau', 1 \rangle$  if and only if  $\tau' = \frac{a\tau + b}{c\tau + d}$ ,  $\begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \text{SL}_2(\mathbb{Z})$

# Lattices

**Lattice** in  $\mathbb{C}$ : a discrete (additive) group of rank 2

**Example:**  $\{a + bi : a, b \in \mathbb{Z}\} = \langle i, 1 \rangle$ .

**Isomorphism of lattices:**  $\Lambda \cong \Lambda'$  if  $\Lambda' = \alpha\Lambda$ ,  $\alpha \in \mathbb{C}$ .

- ▶ Every lattice is isomorphic to  $\langle \tau, 1 \rangle$ ,  $\text{Im } \tau > 0$
- ▶  $\langle \tau, 1 \rangle \cong \langle \tau', 1 \rangle$  if and only if  $\tau' = \frac{a\tau + b}{c\tau + d}$ ,  $\begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \text{SL}_2(\mathbb{Z})$
- ▶  $\{\text{lattices up to isomorphism}\} = \text{SL}_2(\mathbb{Z}) \backslash \mathbb{H}$

$\mathbb{H} = \{\tau \in \mathbb{C} : \text{Im } \tau > 0\}$  “Poincaré (half)plane”

## $j$ -invariant

- ▶  **$j$ -invariant:**  $SL_2(\mathbb{Z})$ -automorphic function on  $\mathbb{H}$  satisfying
$$j(i) = 1728, \quad j\left(\frac{1+\sqrt{-3}}{2}\right) = 0, \quad j(\infty) = \infty$$

# $j$ -invariant

- ▶  **$j$ -invariant:**  $\mathrm{SL}_2(\mathbb{Z})$ -automorphic function on  $\mathbb{H}$  satisfying
$$j(i) = 1728, \quad j\left(\frac{1+\sqrt{-3}}{2}\right) = 0, \quad j(\infty) = \infty$$
- ▶ “ $\mathrm{SL}_2(\mathbb{Z})$ -automorphic” means:  $j(\tau) = j\left(\frac{a\tau+b}{c\tau+d}\right)$ ,
$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathrm{SL}_2(\mathbb{Z})$$

# $j$ -invariant

- ▶  **$j$ -invariant:**  $SL_2(\mathbb{Z})$ -automorphic function on  $\mathbb{H}$  satisfying

$$j(i) = 1728, \quad j\left(\frac{1+\sqrt{-3}}{2}\right) = 0, \quad j(\infty) = \infty$$

- ▶ “ $SL_2(\mathbb{Z})$ -automorphic” means:  $j(\tau) = j\left(\frac{a\tau+b}{c\tau+d}\right)$ ,

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \in SL_2(\mathbb{Z})$$

- ▶ “**Fourier expansion**”

$$j(\tau) = q^{-1} + 744 + 196884q + 21493760q^2 + \dots,$$

$$q = q(\tau) = e^{2\pi i\tau}$$

# $j$ -invariant

- ▶  **$j$ -invariant:**  $\mathrm{SL}_2(\mathbb{Z})$ -automorphic function on  $\mathbb{H}$  satisfying
$$j(i) = 1728, \quad j\left(\frac{1+\sqrt{-3}}{2}\right) = 0, \quad j(\infty) = \infty$$
- ▶ “ $\mathrm{SL}_2(\mathbb{Z})$ -automorphic” means:  $j(\tau) = j\left(\frac{a\tau+b}{c\tau+d}\right)$ ,
$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathrm{SL}_2(\mathbb{Z})$$
- ▶ “**Fourier expansion**”
$$j(\tau) = q^{-1} + 744 + 196884q + 21493760q^2 + \dots,$$
$$q = q(\tau) = e^{2\pi i\tau}$$
- ▶ (remark important in the sequel)  
 $|q|$  small when  $\mathrm{Im} \tau$  large

# $j$ -invariant

- ▶  **$j$ -invariant:**  $\mathrm{SL}_2(\mathbb{Z})$ -automorphic function on  $\mathbb{H}$  satisfying
$$j(i) = 1728, \quad j\left(\frac{1+\sqrt{-3}}{2}\right) = 0, \quad j(\infty) = \infty$$
- ▶ “ $\mathrm{SL}_2(\mathbb{Z})$ -automorphic” means:  $j(\tau) = j\left(\frac{a\tau+b}{c\tau+d}\right)$ ,
$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathrm{SL}_2(\mathbb{Z})$$
- ▶ “**Fourier expansion**”
$$j(\tau) = q^{-1} + 744 + 196884q + 21493760q^2 + \dots,$$
$$q = q(\tau) = e^{2\pi i\tau}$$
- ▶ (remark important in the sequel)  
 $|q|$  small when  $\mathrm{Im} \tau$  large

# $j$ -invariant

- ▶  **$j$ -invariant:**  $\mathrm{SL}_2(\mathbb{Z})$ -automorphic function on  $\mathbb{H}$  satisfying

$$j(i) = 1728, \quad j\left(\frac{1+\sqrt{-3}}{2}\right) = 0, \quad j(\infty) = \infty$$

- ▶ “ $\mathrm{SL}_2(\mathbb{Z})$ -automorphic” means:  $j(\tau) = j\left(\frac{a\tau+b}{c\tau+d}\right)$ ,

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathrm{SL}_2(\mathbb{Z})$$

- ▶ “**Fourier expansion**”

$$j(\tau) = q^{-1} + 744 + 196884q + 21493760q^2 + \dots,$$

$$q = q(\tau) = e^{2\pi i\tau}$$

- ▶ (remark important in the sequel)

$|q|$  small when  $\mathrm{Im} \tau$  large  $\implies |j(\tau)|$  large when  $\mathrm{Im} \tau$  large



# $j$ -invariant

- ▶  **$j$ -invariant:**  $\mathrm{SL}_2(\mathbb{Z})$ -automorphic function on  $\mathbb{H}$  satisfying
$$j(i) = 1728, \quad j\left(\frac{1+\sqrt{-3}}{2}\right) = 0, \quad j(\infty) = \infty$$
- ▶ “ $\mathrm{SL}_2(\mathbb{Z})$ -automorphic” means:  $j(\tau) = j\left(\frac{a\tau+b}{c\tau+d}\right)$ ,
$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathrm{SL}_2(\mathbb{Z})$$
- ▶ **“Fourier expansion”**
$$j(\tau) = q^{-1} + 744 + 196884q + 21493760q^2 + \dots,$$
$$q = q(\tau) = e^{2\pi i\tau}$$
- ▶ (remark important in the sequel)  
 $|q|$  small when  $\mathrm{Im} \tau$  large  $\implies |j(\tau)|$  large when  $\mathrm{Im} \tau$  large
- ▶  $j$ -invariant “classifies lattices”:
$$\langle \tau, 1 \rangle \cong \langle \tau', 1 \rangle \iff j(\tau) = j(\tau')$$

# Complex Multiplication

- ▶  $\text{End}(\Lambda) = \{\alpha \in \mathbb{C} : \alpha\Lambda \subseteq \Lambda\}$ ;

# Complex Multiplication

- ▶  $\text{End}(\Lambda) = \{\alpha \in \mathbb{C} : \alpha\Lambda \subseteq \Lambda\}$ ;

# Complex Multiplication

- ▶  $\text{End}(\Lambda) = \{\alpha \in \mathbb{C} : \alpha\Lambda \subseteq \Lambda\}$ ;       $\text{End}\Lambda \supseteq \mathbb{Z}$
- ▶  $\Lambda$  has **Complex Multiplication** if  $\text{End}\Lambda \not\cong \mathbb{Z}$

# Complex Multiplication

- ▶  $\text{End}(\Lambda) = \{\alpha \in \mathbb{C} : \alpha\Lambda \subseteq \Lambda\}$ ;       $\text{End}\Lambda \supseteq \mathbb{Z}$
- ▶  $\Lambda$  has **Complex Multiplication** if  $\text{End}\Lambda \not\cong \mathbb{Z}$
- ▶  $\Lambda = \langle \tau, 1 \rangle$  has CM  $\iff [\mathbb{Q}(\tau) : \mathbb{Q}] = 2$

# Complex Multiplication

- ▶  $\text{End}(\Lambda) = \{\alpha \in \mathbb{C} : \alpha\Lambda \subseteq \Lambda\}$ ;       $\text{End}\Lambda \supseteq \mathbb{Z}$
- ▶  $\Lambda$  has **Complex Multiplication** if  $\text{End}\Lambda \not\cong \mathbb{Z}$
- ▶  $\Lambda = \langle \tau, 1 \rangle$  has CM  $\iff [\mathbb{Q}(\tau) : \mathbb{Q}] = 2$
- ▶ In this case:

# Complex Multiplication

- ▶  $\text{End}(\Lambda) = \{\alpha \in \mathbb{C} : \alpha\Lambda \subseteq \Lambda\}$ ;       $\text{End}\Lambda \supseteq \mathbb{Z}$
- ▶  $\Lambda$  has **Complex Multiplication** if  $\text{End}\Lambda \not\cong \mathbb{Z}$
- ▶  $\Lambda = \langle \tau, 1 \rangle$  has CM  $\iff [\mathbb{Q}(\tau) : \mathbb{Q}] = 2$
- ▶ In this case:

# Complex Multiplication

- ▶  $\text{End}(\Lambda) = \{\alpha \in \mathbb{C} : \alpha\Lambda \subseteq \Lambda\}$ ;       $\text{End}\Lambda \supseteq \mathbb{Z}$
- ▶  $\Lambda$  has **Complex Multiplication** if  $\text{End}\Lambda \not\cong \mathbb{Z}$
- ▶  $\Lambda = \langle \tau, 1 \rangle$  has CM  $\iff [\mathbb{Q}(\tau) : \mathbb{Q}] = 2$
- ▶ In this case:
  - ▶  $\mathcal{O} = \text{End}\Lambda$  is an order in  $K = \mathbb{Q}(\tau)$  of discriminant  $\Delta = Df^2$ ;



# Complex Multiplication

- ▶  $\text{End}(\Lambda) = \{\alpha \in \mathbb{C} : \alpha\Lambda \subseteq \Lambda\}$ ;       $\text{End}\Lambda \supseteq \mathbb{Z}$
- ▶  $\Lambda$  has **Complex Multiplication** if  $\text{End}\Lambda \not\cong \mathbb{Z}$
- ▶  $\Lambda = \langle \tau, 1 \rangle$  has CM  $\iff [\mathbb{Q}(\tau) : \mathbb{Q}] = 2$
- ▶ In this case:
  - ▶  $\mathcal{O} = \text{End}\Lambda$  is an order in  $K = \mathbb{Q}(\tau)$  of discriminant  $\Delta = Df^2$ ;
  - ▶  $D = D_K < 0$  the **fundamental discriminant**;

# Complex Multiplication

- ▶  $\text{End}(\Lambda) = \{\alpha \in \mathbb{C} : \alpha\Lambda \subseteq \Lambda\}$ ;       $\text{End}\Lambda \supseteq \mathbb{Z}$
- ▶  $\Lambda$  has **Complex Multiplication** if  $\text{End}\Lambda \not\cong \mathbb{Z}$
- ▶  $\Lambda = \langle \tau, 1 \rangle$  has CM  $\iff [\mathbb{Q}(\tau) : \mathbb{Q}] = 2$
- ▶ In this case:
  - ▶  $\mathcal{O} = \text{End}\Lambda$  is an order in  $K = \mathbb{Q}(\tau)$  of discriminant  $\Delta = Df^2$ ;
  - ▶  $D = D_K < 0$  the **fundamental discriminant**;
  - ▶  $f = [\mathcal{O}_K : \mathcal{O}]$  the **conductor**;

# Complex Multiplication

- ▶  $\text{End}(\Lambda) = \{\alpha \in \mathbb{C} : \alpha\Lambda \subseteq \Lambda\}$ ;       $\text{End}\Lambda \supseteq \mathbb{Z}$
- ▶  $\Lambda$  has **Complex Multiplication** if  $\text{End}\Lambda \not\subseteq \mathbb{Z}$
- ▶  $\Lambda = \langle \tau, 1 \rangle$  has CM  $\iff [\mathbb{Q}(\tau) : \mathbb{Q}] = 2$
- ▶ In this case:
  - ▶  $\mathcal{O} = \text{End}\Lambda$  is an order in  $K = \mathbb{Q}(\tau)$  of discriminant  $\Delta = Df^2$ ;
  - ▶  $D = D_K < 0$  the **fundamental discriminant**;
  - ▶  $f = [\mathcal{O}_K : \mathcal{O}]$  the **conductor**;
  - ▶  $\mathcal{O} = \mathbb{Z} + f\mathcal{O}_K = \mathbb{Z} \left[ \frac{\Delta + \sqrt{\Delta}}{2} \right] =: \mathcal{O}_\Delta$ ;

# Complex Multiplication

- ▶  $\text{End}(\Lambda) = \{\alpha \in \mathbb{C} : \alpha\Lambda \subseteq \Lambda\}; \quad \text{End}\Lambda \supseteq \mathbb{Z}$
- ▶  $\Lambda$  has **Complex Multiplication** if  $\text{End}\Lambda \not\subseteq \mathbb{Z}$
- ▶  $\Lambda = \langle \tau, 1 \rangle$  has CM  $\iff [\mathbb{Q}(\tau) : \mathbb{Q}] = 2$
- ▶ In this case:
  - ▶  $\mathcal{O} = \text{End}\Lambda$  is an order in  $K = \mathbb{Q}(\tau)$  of discriminant  $\Delta = Df^2$ ;
  - ▶  $D = D_K < 0$  the **fundamental discriminant**;
  - ▶  $f = [\mathcal{O}_K : \mathcal{O}]$  the **conductor**;
  - ▶  $\mathcal{O} = \mathbb{Z} + f\mathcal{O}_K = \mathbb{Z} \left[ \frac{\Delta + \sqrt{\Delta}}{2} \right] =: \mathcal{O}_\Delta$ ;
  - ▶ if  $\tau$  is root of  $at^2 + bt + c \in \mathbb{Z}[t]$ ,

# Complex Multiplication

- ▶  $\text{End}(\Lambda) = \{\alpha \in \mathbb{C} : \alpha\Lambda \subseteq \Lambda\}; \quad \text{End}\Lambda \supseteq \mathbb{Z}$
- ▶  $\Lambda$  has **Complex Multiplication** if  $\text{End}\Lambda \not\subseteq \mathbb{Z}$
- ▶  $\Lambda = \langle \tau, 1 \rangle$  has CM  $\iff [\mathbb{Q}(\tau) : \mathbb{Q}] = 2$
- ▶ In this case:
  - ▶  $\mathcal{O} = \text{End}\Lambda$  is an order in  $K = \mathbb{Q}(\tau)$  of discriminant  $\Delta = Df^2$ ;
  - ▶  $D = D_K < 0$  the **fundamental discriminant**;
  - ▶  $f = [\mathcal{O}_K : \mathcal{O}]$  the **conductor**;
  - ▶  $\mathcal{O} = \mathbb{Z} + f\mathcal{O}_K = \mathbb{Z} \left[ \frac{\Delta + \sqrt{\Delta}}{2} \right] =: \mathcal{O}_\Delta$ ;
  - ▶ if  $\tau$  is root of  $at^2 + bt + c \in \mathbb{Z}[t]$ ,

# Complex Multiplication

- ▶  $\text{End}(\Lambda) = \{\alpha \in \mathbb{C} : \alpha\Lambda \subseteq \Lambda\}; \quad \text{End}\Lambda \supseteq \mathbb{Z}$
- ▶  $\Lambda$  has **Complex Multiplication** if  $\text{End}\Lambda \not\subseteq \mathbb{Z}$
- ▶  $\Lambda = \langle \tau, 1 \rangle$  has CM  $\iff [\mathbb{Q}(\tau) : \mathbb{Q}] = 2$
- ▶ In this case:
  - ▶  $\mathcal{O} = \text{End}\Lambda$  is an order in  $K = \mathbb{Q}(\tau)$  of discriminant  $\Delta = Df^2$ ;
  - ▶  $D = D_K < 0$  the **fundamental discriminant**;
  - ▶  $f = [\mathcal{O}_K : \mathcal{O}]$  the **conductor**;
  - ▶  $\mathcal{O} = \mathbb{Z} + f\mathcal{O}_K = \mathbb{Z} \left[ \frac{\Delta + \sqrt{\Delta}}{2} \right] =: \mathcal{O}_\Delta$ ;
  - ▶ if  $\tau$  is root of  $at^2 + bt + c \in \mathbb{Z}[t]$ ,  $(a, b, c) = 1$

# Complex Multiplication

- ▶  $\text{End}(\Lambda) = \{\alpha \in \mathbb{C} : \alpha\Lambda \subseteq \Lambda\}$ ;       $\text{End}\Lambda \supseteq \mathbb{Z}$
- ▶  $\Lambda$  has **Complex Multiplication** if  $\text{End}\Lambda \not\cong \mathbb{Z}$
- ▶  $\Lambda = \langle \tau, 1 \rangle$  has CM  $\iff [\mathbb{Q}(\tau) : \mathbb{Q}] = 2$
- ▶ In this case:
  - ▶  $\mathcal{O} = \text{End}\Lambda$  is an order in  $K = \mathbb{Q}(\tau)$  of discriminant  $\Delta = Df^2$ ;
  - ▶  $D = D_K < 0$  the **fundamental discriminant**;
  - ▶  $f = [\mathcal{O}_K : \mathcal{O}]$  the **conductor**;
  - ▶  $\mathcal{O} = \mathbb{Z} + f\mathcal{O}_K = \mathbb{Z} \left[ \frac{\Delta + \sqrt{\Delta}}{2} \right] =: \mathcal{O}_\Delta$ ;
  - ▶ if  $\tau$  is root of  $at^2 + bt + c \in \mathbb{Z}[t]$ ,  $(a, b, c) = 1$  then  $\Delta = b^2 - 4ac$  and  $\tau = \frac{-b + \sqrt{\Delta}}{2a}$ .

# Class Field Theory

- ▶  $j(\tau)$  algebraic number (even algebraic integer)



# Class Field Theory

- ▶  $j(\tau)$  algebraic number (even algebraic integer)
- ▶  $K(j(\tau))$  is abelian extension of  $K = \mathbb{Q}(\tau)$  (the “Ring Class Field”)

# Class Field Theory

- ▶  $j(\tau)$  algebraic number (even algebraic integer)
- ▶  $K(j(\tau))$  is abelian extension of  $K = \mathbb{Q}(\tau)$  (the “Ring Class Field”)
- ▶  $[K(j(\tau)) : K] = [\mathbb{Q}(j(\tau)) : \mathbb{Q}] = h(\Delta)$

# Class Field Theory

- ▶  $j(\tau)$  algebraic number (even algebraic integer)
- ▶  $K(j(\tau))$  is abelian extension of  $K = \mathbb{Q}(\tau)$  (the “Ring Class Field”)
- ▶  $[K(j(\tau)) : K] = [\mathbb{Q}(j(\tau)) : \mathbb{Q}] = h(\Delta)$
- ▶  $h(\Delta)$  the class number of the order  $\mathcal{O}_\Delta$

# Class Field Theory

- ▶  $j(\tau)$  algebraic number (even algebraic integer)
- ▶  $K(j(\tau))$  is abelian extension of  $K = \mathbb{Q}(\tau)$  (the “Ring Class Field”)
- ▶  $[K(j(\tau)) : K] = [\mathbb{Q}(j(\tau)) : \mathbb{Q}] = h(\Delta)$
- ▶  $h(\Delta)$  the class number of the order  $\mathcal{O}_\Delta$
- ▶ moreover:  $\text{Gal}(K(j(\tau))/K) = \text{Cl}(\Delta)$

# The Class Number

- ▶  $h(\Delta) \rightarrow \infty$  as  $|\Delta| \rightarrow \infty$  (Siegel)

# The Class Number

- ▶  $h(\Delta) \rightarrow \infty$  as  $|\Delta| \rightarrow \infty$  (Siegel)
- ▶ In other words, for  $h \in \mathbb{Z}_{>0}$  there exist finitely many  $\Delta$  with  $h(\Delta) = h$ .

# The Class Number

- ▶  $h(\Delta) \rightarrow \infty$  as  $|\Delta| \rightarrow \infty$  (Siegel)
- ▶ In other words, for  $h \in \mathbb{Z}_{>0}$  there exist finitely many  $\Delta$  with  $h(\Delta) = h$ .
- ▶ In particular (Heegner-Stark) there exist thirteen  $\Delta$  with  $h(\Delta) = 1$  (the corresponding  $j$  belong to  $\mathbb{Z}$ ):

$\Delta$	$-3$	$-3 \cdot 2^2$	$-3 \cdot 3^3$	$-4$	$-4 \cdot 2^2$	$-7$	$-7 \cdot 2^2$	$-8$
$j$	$0$	$2^4 3^3 5^3$	$-2^{15} 3 \cdot 5^3$	$2^6 3^3$	$2^3 3^3 11^3$	$-3^3 5^3$	$3^3 5^3 17^3$	$2^6 5^3$
$\Delta$	$-11$	$-19$	$-43$	$-67$	$-163$			
$j$	$-2^{15}$	$-2^{15} 3^3$	$-2^{18} 3^3 5^3$	$-2^{15} 3^3 5^3 11^3$	$-2^{18} 3^3 5^3 23^3 29^3$			

# The Class Number

- ▶  $h(\Delta) \rightarrow \infty$  as  $|\Delta| \rightarrow \infty$  (Siegel)
- ▶ In other words, for  $h \in \mathbb{Z}_{>0}$  there exist finitely many  $\Delta$  with  $h(\Delta) = h$ .
- ▶ In particular (Heegner-Stark) there exist thirteen  $\Delta$  with  $h(\Delta) = 1$  (the corresponding  $j$  belong to  $\mathbb{Z}$ ):

$\Delta$	$-3$	$-3 \cdot 2^2$	$-3 \cdot 3^3$	$-4$	$-4 \cdot 2^2$	$-7$	$-7 \cdot 2^2$	$-8$
$j$	$0$	$2^4 3^3 5^3$	$-2^{15} 3 \cdot 5^3$	$2^6 3^3$	$2^3 3^3 11^3$	$-3^3 5^3$	$3^3 5^3 17^3$	$2^6 5^3$
$\Delta$	$-11$	$-19$	$-43$	$-67$	$-163$			
$j$	$-2^{15}$	$-2^{15} 3^3$	$-2^{18} 3^3 5^3$	$-2^{15} 3^3 5^3 11^3$	$-2^{18} 3^3 5^3 23^3 29^3$			

- ▶ A funny example (Hermite):

$$e^{\pi\sqrt{163}} = 262537412640768743.99999999999925007 \dots$$



# The Class Number

- ▶  $h(\Delta) \rightarrow \infty$  as  $|\Delta| \rightarrow \infty$  (Siegel)
- ▶ In other words, for  $h \in \mathbb{Z}_{>0}$  there exist finitely many  $\Delta$  with  $h(\Delta) = h$ .
- ▶ In particular (Heegner-Stark) there exist thirteen  $\Delta$  with  $h(\Delta) = 1$  (the corresponding  $j$  belong to  $\mathbb{Z}$ ):

$\Delta$	$-3$	$-3 \cdot 2^2$	$-3 \cdot 3^3$	$-4$	$-4 \cdot 2^2$	$-7$	$-7 \cdot 2^2$	$-8$
$j$	$0$	$2^4 3^3 5^3$	$-2^{15} 3 \cdot 5^3$	$2^6 3^3$	$2^3 3^3 11^3$	$-3^3 5^3$	$3^3 5^3 17^3$	$2^6 5^3$
$\Delta$	$-11$	$-19$	$-43$	$-67$	$-163$			
$j$	$-2^{15}$	$-2^{15} 3^3$	$-2^{18} 3^3 5^3$	$-2^{15} 3^3 5^3 11^3$	$-2^{18} 3^3 5^3 23^3 29^3$			

- ▶ A funny example (Hermite):

$$e^{\pi\sqrt{163}} = 262537412640768743.99999999999925007 \dots$$

# The Class Number

- ▶  $h(\Delta) \rightarrow \infty$  as  $|\Delta| \rightarrow \infty$  (Siegel)
- ▶ In other words, for  $h \in \mathbb{Z}_{>0}$  there exist finitely many  $\Delta$  with  $h(\Delta) = h$ .
- ▶ In particular (Heegner-Stark) there exist thirteen  $\Delta$  with  $h(\Delta) = 1$  (the corresponding  $j$  belong to  $\mathbb{Z}$ ):

$$\begin{array}{l|ccccccc} \Delta & -3 & -3 \cdot 2^2 & -3 \cdot 3^3 & -4 & -4 \cdot 2^2 & -7 & -7 \cdot 2^2 & -8 \\ j & 0 & 2^4 3^3 5^3 & -2^{15} 3 \cdot 5^3 & 2^6 3^3 & 2^3 3^3 11^3 & -3^3 5^3 & 3^3 5^3 17^3 & 2^6 5^3 \\ \hline \Delta & -11 & -19 & -43 & -67 & -163 \\ j & -2^{15} & -2^{15} 3^3 & -2^{18} 3^3 5^3 & -2^{15} 3^3 5^3 11^3 & -2^{18} 3^3 5^3 23^3 29^3 \end{array}$$

- ▶ A funny example (Hermite):

$$e^{\pi\sqrt{163}} = 262537412640768743.99999999999925007 \dots$$

- ▶  $\tau = \frac{1+\sqrt{-163}}{2}$

# The Class Number

- ▶  $h(\Delta) \rightarrow \infty$  as  $|\Delta| \rightarrow \infty$  (Siegel)
- ▶ In other words, for  $h \in \mathbb{Z}_{>0}$  there exist finitely many  $\Delta$  with  $h(\Delta) = h$ .
- ▶ In particular (Heegner-Stark) there exist thirteen  $\Delta$  with  $h(\Delta) = 1$  (the corresponding  $j$  belong to  $\mathbb{Z}$ ):

$\Delta$	$-3$	$-3 \cdot 2^2$	$-3 \cdot 3^3$	$-4$	$-4 \cdot 2^2$	$-7$	$-7 \cdot 2^2$	$-8$
$j$	$0$	$2^4 3^3 5^3$	$-2^{15} 3 \cdot 5^3$	$2^6 3^3$	$2^3 3^3 11^3$	$-3^3 5^3$	$3^3 5^3 17^3$	$2^6 5^3$
$\Delta$	$-11$	$-19$	$-43$	$-67$	$-163$			
$j$	$-2^{15}$	$-2^{15} 3^3$	$-2^{18} 3^3 5^3$	$-2^{15} 3^3 5^3 11^3$	$-2^{18} 3^3 5^3 23^3 29^3$			

- ▶ A funny example (Hermite):

$$e^{\pi\sqrt{163}} = 262537412640768743.99999999999925007 \dots$$

- ▶  $\tau = \frac{1+\sqrt{-163}}{2}$
- ▶  $e^{\pi\sqrt{163}} = -e^{2\pi i\tau} \approx -j(\tau) + 744 \in \mathbb{Z}$

# The Class Number

- ▶  $h(\Delta) \rightarrow \infty$  as  $|\Delta| \rightarrow \infty$  (Siegel)
- ▶ In other words, for  $h \in \mathbb{Z}_{>0}$  there exist finitely many  $\Delta$  with  $h(\Delta) = h$ .
- ▶ In particular (Heegner-Stark) there exist thirteen  $\Delta$  with  $h(\Delta) = 1$  (the corresponding  $j$  belong to  $\mathbb{Z}$ ):

$$\begin{array}{l|cccccccc} \Delta & -3 & -3 \cdot 2^2 & -3 \cdot 3^3 & -4 & -4 \cdot 2^2 & -7 & -7 \cdot 2^2 & -8 \\ j & 0 & 2^4 3^3 5^3 & -2^{15} 3 \cdot 5^3 & 2^6 3^3 & 2^3 3^3 11^3 & -3^3 5^3 & 3^3 5^3 17^3 & 2^6 5^3 \\ \hline \Delta & -11 & -19 & -43 & -67 & -163 & & & \\ j & -2^{15} & -2^{15} 3^3 & -2^{18} 3^3 5^3 & -2^{15} 3^3 5^3 11^3 & -2^{18} 3^3 5^3 23^3 29^3 & & & \end{array}$$

- ▶ A funny example (Hermite):

$$e^{\pi\sqrt{163}} = 262537412640768743.999999999999925007 \dots$$

- ▶  $\tau = \frac{1+\sqrt{-163}}{2}$

- ▶  $e^{\pi\sqrt{163}} = -e^{2\pi i\tau} \approx -j(\tau) + 744 \in \mathbb{Z}$

- ▶ Currently all  $\Delta$  with  $h_{\Delta} \leq 100$  are known (Watkins 2006).

## Complex Multiplication

Lattices

$j$ -invariant

Complex Multiplication

Class Field Theory

The Class Number

## Theorem of André

Special Points and Special Curves

Theorem of André

## CM-Points on Straight Lines

Kühne's "uniformity observation"

CM-Points on Straight Lines

## The Proof

Equality of CM-fields

The Proof

## Proof of Theorem ECMF

Discriminants with Class Group Annihilated by 2

Proof of Theorem ECMF

# Special Points and Special Curves

$\tau$  imaginary quadratic  $\Rightarrow j(\tau) \in \bar{\mathbb{Q}}$

# Special Points and Special Curves

$\tau$  imaginary quadratic  $\Rightarrow j(\tau) \in \bar{\mathbb{Q}}$

**CM-point** or **special point** on  $\mathbb{C}^2$ :  $(j(\tau_1), j(\tau_2))$ .

# Special Points and Special Curves

$\tau$  imaginary quadratic  $\Rightarrow j(\tau) \in \bar{\mathbb{Q}}$

**CM-point** or **special point** on  $\mathbb{C}^2$ :  $(j(\tau_1), j(\tau_2))$ .

**Question:** can an irreducible plane curve  $F(x_1, x_2) = 0$  contain infinitely many CM-points?



# Special Points and Special Curves

$\tau$  imaginary quadratic  $\Rightarrow j(\tau) \in \bar{\mathbb{Q}}$

**CM-point** or **special point** on  $\mathbb{C}^2$ :  $(j(\tau_1), j(\tau_2))$ .

**Question:** can an irreducible plane curve  $F(x_1, x_2) = 0$  contain infinitely many CM-points?

**Special curves:**

- ▶ vertical line  $x_1 = j(\tau_1)$

# Special Points and Special Curves

$\tau$  imaginary quadratic  $\Rightarrow j(\tau) \in \bar{\mathbb{Q}}$

**CM-point** or **special point** on  $\mathbb{C}^2$ :  $(j(\tau_1), j(\tau_2))$ .

**Question:** can an irreducible plane curve  $F(x_1, x_2) = 0$  contain infinitely many CM-points?

**Special curves:**

- ▶ vertical line  $x_1 = j(\tau_1)$
- ▶ horizontal line  $x_2 = j(\tau_2)$

# Special Points and Special Curves

$\tau$  imaginary quadratic  $\Rightarrow j(\tau) \in \bar{\mathbb{Q}}$

**CM-point** or **special point** on  $\mathbb{C}^2$ :  $(j(\tau_1), j(\tau_2))$ .

**Question:** can an irreducible plane curve  $F(x_1, x_2) = 0$  contain infinitely many CM-points?

**Special curves:**

- ▶ vertical line  $x_1 = j(\tau_1)$
- ▶ horizontal line  $x_2 = j(\tau_2)$
- ▶  $Y_0(N)$  realized as  $\Phi_N(x_1, x_2) = 0$

# Modular Curves and Modular Polynomials

- ▶  $\Phi_N(x_1, x_2)$   $N$ th “modular polynomial”:  $\Phi_N(j(z), j(Nz)) = 0$

# Modular Curves and Modular Polynomials

- ▶  $\Phi_N(x_1, x_2)$   $N$ th “modular polynomial”:  $\Phi_N(j(z), j(Nz)) = 0$
- ▶  $(j(\tau), j(N\tau)) \in Y_0(N)$  for every  $\tau$ .

# Modular Curves and Modular Polynomials

- ▶  $\Phi_N(x_1, x_2)$   $N$ th “modular polynomial”:  $\Phi_N(j(z), j(Nz)) = 0$
- ▶  $(j(\tau), j(N\tau)) \in Y_0(N)$  for every  $\tau$ .
- ▶ More generally:  
for  $\gamma \in \mathrm{GL}_2(\mathbb{Q})$  there exists  $N$  such that  
 $(j(\tau), j(\gamma\tau)) \in Y_0(N)$  for every  $\tau$ .

# Modular Curves and Modular Polynomials

- ▶  $\Phi_N(x_1, x_2)$   $N$ th “modular polynomial”:  $\Phi_N(j(z), j(Nz)) = 0$
- ▶  $(j(\tau), j(N\tau)) \in Y_0(N)$  for every  $\tau$ .
- ▶ More generally:  
for  $\gamma \in \mathrm{GL}_2(\mathbb{Q})$  there exists  $N$  such that  
 $(j(\tau), j(\gamma\tau)) \in Y_0(N)$  for every  $\tau$ .

# Modular Curves and Modular Polynomials

- ▶  $\Phi_N(x_1, x_2)$   $N$ th “modular polynomial”:  $\Phi_N(j(z), j(Nz)) = 0$
- ▶  $(j(\tau), j(N\tau)) \in Y_0(N)$  for every  $\tau$ .
- ▶ More generally:  
for  $\gamma \in \text{GL}_2(\mathbb{Q})$  there exists  $N$  such that  
 $(j(\tau), j(\gamma\tau)) \in Y_0(N)$  for every  $\tau$ .

## Polynomials $\Phi_N, N \leq 3$

$$\Phi_1(x, y) = x - y$$

$$\begin{aligned}\Phi_2(x, y) = & -x^2y^2 + x^3 + y^3 + 1488x^2y + 1488xy^2 + 40773375xy \\ & - 162000x^2 - 162000y^2 + 8748000000x + 8748000000y - 15746400000000\end{aligned}$$

$$\begin{aligned}\Phi_3(x, y) = & x^4 + y^4 - x^3y^3 + 2232x^3y^2 + 2232x^2y^3 - 1069956x^3y - 1069956xy^3 \\ & + 36864000x^3 + 36864000y^3 + 2587918086x^2y^2 \\ & + 8900222976000x^2y + 8900222976000xy^2 + 452984832000000x^2 + 452984832000000y^2 \\ & - 770845966336000000xy + 1855425871872000000000x + 1855425871872000000000y\end{aligned}$$



# Theorem of André

**Theorem** (André, 1998) A non-special irreducible plane curve can have only finitely many special points.

# Theorem of André

**Theorem** (André, 1998) A non-special irreducible plane curve can have only finitely many special points.

**Different proofs:**

# Theorem of André

**Theorem** (André, 1998) A non-special irreducible plane curve can have only finitely many special points.

## Different proofs:

- ▶ André (1998)

# Theorem of André

**Theorem** (André, 1998) A non-special irreducible plane curve can have only finitely many special points.

## Different proofs:

- ▶ André (1998)
- ▶ Edixhoven (1998, GRH)

# Theorem of André

**Theorem** (André, 1998) A non-special irreducible plane curve can have only finitely many special points.

## Different proofs:

- ▶ André (1998)
- ▶ Edixhoven (1998, GRH)
- ▶ Pila (2009, extends to higher dimension)

# Theorem of André

**Theorem** (André, 1998) A non-special irreducible plane curve can have only finitely many special points.

## Different proofs:

- ▶ André (1998)
- ▶ Edixhoven (1998, GRH)
- ▶ Pila (2009, extends to higher dimension)

**All non-effective, use Siegel-Brauer**

# Theorem of André

**Theorem** (André, 1998) A non-special irreducible plane curve can have only finitely many special points.

## Different proofs:

- ▶ André (1998)
- ▶ Edixhoven (1998, GRH)
- ▶ Pila (2009, extends to higher dimension)

**All non-effective, use Siegel-Brauer**

- ▶ Breuer (2001, GRH, effective)

# Theorem of André

**Theorem** (André, 1998) A non-special irreducible plane curve can have only finitely many special points.

## Different proofs:

- ▶ André (1998)
- ▶ Edixhoven (1998, GRH)
- ▶ Pila (2009, extends to higher dimension)

### **All non-effective, use Siegel-Brauer**

- ▶ Breuer (2001, GRH, effective)
- ▶ B., Masser, Zannier (2013, effective)



# Theorem of André

**Theorem** (André, 1998) A non-special irreducible plane curve can have only finitely many special points.

## Different proofs:

- ▶ André (1998)
- ▶ Edixhoven (1998, GRH)
- ▶ Pila (2009, extends to higher dimension)

### **All non-effective, use Siegel-Brauer**

- ▶ Breuer (2001, GRH, effective)
- ▶ B., Masser, Zannier (2013, effective)
- ▶ Kühne (2012, 2013, effective)

# Theorem of André

**Theorem** (André, 1998) A non-special irreducible plane curve can have only finitely many special points.

## Different proofs:

- ▶ André (1998)
- ▶ Edixhoven (1998, GRH)
- ▶ Pila (2009, extends to higher dimension)

### **All non-effective, use Siegel-Brauer**

- ▶ Breuer (2001, GRH, effective)
- ▶ B., Masser, Zannier (2013, effective)
- ▶ Kühne (2012, 2013, effective)

# Theorem of André

**Theorem** (André, 1998) A non-special irreducible plane curve can have only finitely many special points.

## Different proofs:

- ▶ André (1998)
- ▶ Edixhoven (1998, GRH)
- ▶ Pila (2009, extends to higher dimension)

### **All non-effective, use Siegel-Brauer**

- ▶ Breuer (2001, GRH, effective)
- ▶ B., Masser, Zannier (2013, effective)
- ▶ Kühne (2012, 2013, effective)

## Particular cases:

# Theorem of André

**Theorem** (André, 1998) A non-special irreducible plane curve can have only finitely many special points.

## Different proofs:

- ▶ André (1998)
- ▶ Edixhoven (1998, GRH)
- ▶ Pila (2009, extends to higher dimension)

### **All non-effective, use Siegel-Brauer**

- ▶ Breuer (2001, GRH, effective)
- ▶ B., Masser, Zannier (2013, effective)
- ▶ Kühne (2012, 2013, effective)

## Particular cases:

- ▶ no CM-points on  $x_1 + x_2 = 1$  (Kühne 2013)

# Theorem of André

**Theorem** (André, 1998) A non-special irreducible plane curve can have only finitely many special points.

## Different proofs:

- ▶ André (1998)
- ▶ Edixhoven (1998, GRH)
- ▶ Pila (2009, extends to higher dimension)

### **All non-effective, use Siegel-Brauer**

- ▶ Breuer (2001, GRH, effective)
- ▶ B., Masser, Zannier (2013, effective)
- ▶ Kühne (2012, 2013, effective)

## Particular cases:

- ▶ no CM-points on  $x_1 + x_2 = 1$  (Kühne 2013)
- ▶ no CM-points on  $x_1 x_2 = 1$  (B., Masser, Zannier 2013)

## Complex Multiplication

Lattices

$j$ -invariant

Complex Multiplication

Class Field Theory

The Class Number

## Theorem of André

Special Points and Special Curves

Theorem of André

## CM-Points on Straight Lines

Kühne's "uniformity observation"

CM-Points on Straight Lines

## The Proof

Equality of CM-fields

The Proof

## Proof of Theorem ECMF

Discriminants with Class Group Annihilated by 2

Proof of Theorem ECMF

# Kühne's "uniformity observation"

## Kühne (2013):

- ▶ If  $(j(\tau_1), j(\tau_2))$  belongs to a non-special straight line over a n.f.  $L$ , then  $|\Delta_1|, |\Delta_2| \leq c_{\text{eff}}([L : \mathbb{Q}])$ .

# Kühne's "uniformity observation"

## Kühne (2013):

- ▶ If  $(j(\tau_1), j(\tau_2))$  belongs to a non-special straight line over a n.f.  $L$ , then  $|\Delta_1|, |\Delta_2| \leq c_{\text{eff}}([L : \mathbb{Q}])$ .  
( $\Delta_i$  discriminant of the CM-order  $\text{End}\langle \tau_i, 1 \rangle$ )



# Kühne's "uniformity observation"

## Kühne (2013):

- ▶ If  $(j(\tau_1), j(\tau_2))$  belongs to a non-special straight line over a n.f.  $L$ , then  $|\Delta_1|, |\Delta_2| \leq c_{\text{eff}}([L : \mathbb{Q}])$ .  
( $\Delta_j$  discriminant of the CM-order  $\text{End}\langle \tau_j, 1 \rangle$ )
- ▶ In particular: all CM-points belonging to non-special straight lines defined over  $\mathbb{Q}$  can (in principle) be listed explicitly.

# Kühne's "uniformity observation"

## Kühne (2013):

- ▶ If  $(j(\tau_1), j(\tau_2))$  belongs to a non-special straight line over a n.f.  $L$ , then  $|\Delta_1|, |\Delta_2| \leq c_{\text{eff}}([L : \mathbb{Q}])$ .  
( $\Delta_i$  discriminant of the CM-order  $\text{End}\langle \tau_i, 1 \rangle$ )
- ▶ In particular: all CM-points belonging to non-special straight lines defined over  $\mathbb{Q}$  can (in principle) be listed explicitly.
- ▶ Bajolet (2014): software to determine all CM-points on a given line.

# CM-Points on Straight Lines

**Special straight lines:**

# CM-Points on Straight Lines

## Special straight lines:

- ▶ vertical  $x_1 = j(\tau_1)$  and horizontal  $x_2 = j(\tau_2)$  lines;

# CM-Points on Straight Lines

## Special straight lines:

- ▶ vertical  $x_1 = j(\tau_1)$  and horizontal  $x_2 = j(\tau_2)$  lines;
- ▶  $x_1 = x_2$  (which is  $Y_0(1)$ ).

# CM-Points on Straight Lines

## Special straight lines:

- ▶ vertical  $x_1 = j(\tau_1)$  and horizontal  $x_2 = j(\tau_2)$  lines;
- ▶  $x_1 = x_2$  (which is  $Y_0(1)$ ).

# CM-Points on Straight Lines

## Special straight lines:

- ▶ vertical  $x_1 = j(\tau_1)$  and horizontal  $x_2 = j(\tau_2)$  lines;
- ▶  $x_1 = x_2$  (which is  $Y_0(1)$ ).

**Obvious cases**  $(j(\tau_1), j(\tau_2))$  belongs to a non-special straight line over  $\mathbb{Q}$  in one of the following cases:

# CM-Points on Straight Lines

## Special straight lines:

- ▶ vertical  $x_1 = j(\tau_1)$  and horizontal  $x_2 = j(\tau_2)$  lines;
- ▶  $x_1 = x_2$  (which is  $Y_0(1)$ ).

**Obvious cases**  $(j(\tau_1), j(\tau_2))$  belongs to a non-special straight line over  $\mathbb{Q}$  in one of the following cases:

- ▶  $j(\tau_1), j(\tau_2) \in \mathbb{Q}$ ;



# CM-Points on Straight Lines

## Special straight lines:

- ▶ vertical  $x_1 = j(\tau_1)$  and horizontal  $x_2 = j(\tau_2)$  lines;
- ▶  $x_1 = x_2$  (which is  $Y_0(1)$ ).

**Obvious cases**  $(j(\tau_1), j(\tau_2))$  belongs to a non-special straight line over  $\mathbb{Q}$  in one of the following cases:

- ▶  $j(\tau_1), j(\tau_2) \in \mathbb{Q}$ ;
- ▶  $j(\tau_1) \neq j(\tau_2)$ ,  $\mathbb{Q}(j(\tau_1)) = \mathbb{Q}(j(\tau_2)) = K$ ,  $[K : \mathbb{Q}] = 2$ .

# CM-Points on Straight Lines

## Special straight lines:

- ▶ vertical  $x_1 = j(\tau_1)$  and horizontal  $x_2 = j(\tau_2)$  lines;
- ▶  $x_1 = x_2$  (which is  $Y_0(1)$ ).

**Obvious cases**  $(j(\tau_1), j(\tau_2))$  belongs to a non-special straight line over  $\mathbb{Q}$  in one of the following cases:

- ▶  $j(\tau_1), j(\tau_2) \in \mathbb{Q}$ ;
- ▶  $j(\tau_1) \neq j(\tau_2)$ ,  $\mathbb{Q}(j(\tau_1)) = \mathbb{Q}(j(\tau_2)) = K$ ,  $[K : \mathbb{Q}] = 2$ .

# CM-Points on Straight Lines

## Special straight lines:

- ▶ vertical  $x_1 = j(\tau_1)$  and horizontal  $x_2 = j(\tau_2)$  lines;
- ▶  $x_1 = x_2$  (which is  $Y_0(1)$ ).

**Obvious cases**  $(j(\tau_1), j(\tau_2))$  belongs to a non-special straight line over  $\mathbb{Q}$  in one of the following cases:

- ▶  $j(\tau_1), j(\tau_2) \in \mathbb{Q}$ ;
- ▶  $j(\tau_1) \neq j(\tau_2)$ ,  $\mathbb{Q}(j(\tau_1)) = \mathbb{Q}(j(\tau_2)) = K$ ,  $[K : \mathbb{Q}] = 2$ .

(Can be easily listed.)

# CM-Points on Straight Lines

## Special straight lines:

- ▶ vertical  $x_1 = j(\tau_1)$  and horizontal  $x_2 = j(\tau_2)$  lines;
- ▶  $x_1 = x_2$  (which is  $Y_0(1)$ ).

**Obvious cases**  $(j(\tau_1), j(\tau_2))$  belongs to a non-special straight line over  $\mathbb{Q}$  in one of the following cases:

- ▶  $j(\tau_1), j(\tau_2) \in \mathbb{Q}$ ;
- ▶  $j(\tau_1) \neq j(\tau_2)$ ,  $\mathbb{Q}(j(\tau_1)) = \mathbb{Q}(j(\tau_2)) = K$ ,  $[K : \mathbb{Q}] = 2$ .

(Can be easily listed.)

**Theorem** (A., B., Pizarro; May 2014) If a CM-points belongs to a non-special straight line over  $\mathbb{Q}$  then we have one of the two cases above.

## Complex Multiplication

Lattices

$j$ -invariant

Complex Multiplication

Class Field Theory

The Class Number

## Theorem of André

Special Points and Special Curves

Theorem of André

## CM-Points on Straight Lines

Kühne's "uniformity observation"

CM-Points on Straight Lines

## The Proof

Equality of CM-fields

The Proof

## Proof of Theorem ECMF

Discriminants with Class Group Annihilated by 2

Proof of Theorem ECMF

# Equality of CM-fields

**Theorem ECMF** (based on ideas of André, Edixhoven and Kühne) Assume that  $L = \mathbb{Q}(j(\tau_1)) = \mathbb{Q}(j(\tau_2))$ .

# Equality of CM-fields

**Theorem ECMF** (based on ideas of André, Edixhoven and Kühne) Assume that  $L = \mathbb{Q}(j(\tau_1)) = \mathbb{Q}(j(\tau_2))$ .

- If  $\mathbb{Q}(\tau_1) \neq \mathbb{Q}(\tau_2)$  then  $L$  is the table:

Field $L$	$\Delta$	$\text{Cl}(\Delta)$
$\mathbb{Q}$	$-3, -4, -7, -8, -11, -12, -16, -19, -27, -28, -43, -67, -163$	trivial
$\mathbb{Q}(\sqrt{2})$	$-24, -32, -64, -88$	$\mathbb{Z}/2\mathbb{Z}$
$\mathbb{Q}(\sqrt{3})$	$-36, -48$	$\mathbb{Z}/2\mathbb{Z}$
$\mathbb{Q}(\sqrt{5})$	$-15, -20, -35, -40, -60, -75, -100, -115, -235$	$\mathbb{Z}/2\mathbb{Z}$
$\mathbb{Q}(\sqrt{13})$	$-52, -91, -403$	$\mathbb{Z}/2\mathbb{Z}$
$\mathbb{Q}(\sqrt{17})$	$-51, -187$	$\mathbb{Z}/2\mathbb{Z}$
$\mathbb{Q}(\sqrt{2}, \sqrt{3})$	$-96, -192, -288$	$(\mathbb{Z}/2\mathbb{Z})^2$
$\mathbb{Q}(\sqrt{3}, \sqrt{5})$	$-180, -240$	$(\mathbb{Z}/2\mathbb{Z})^2$
$\mathbb{Q}(\sqrt{5}, \sqrt{13})$	$-195, -520, -715$	$(\mathbb{Z}/2\mathbb{Z})^2$
$\mathbb{Q}(\sqrt{2}, \sqrt{5})$	$-120, -160, -280, -760$	$(\mathbb{Z}/2\mathbb{Z})^2$
$\mathbb{Q}(\sqrt{5}, \sqrt{17})$	$-340, -595$	$(\mathbb{Z}/2\mathbb{Z})^2$
$\mathbb{Q}(\sqrt{2}, \sqrt{3}, \sqrt{5})$	$-480, -960$	$(\mathbb{Z}/2\mathbb{Z})^3$

# Equality of CM-fields

**Theorem ECMF** (based on ideas of André, Edixhoven and Kühne) Assume that  $L = \mathbb{Q}(j(\tau_1)) = \mathbb{Q}(j(\tau_2))$ .

- ▶ If  $\mathbb{Q}(\tau_1) \neq \mathbb{Q}(\tau_2)$  then  $L$  is the table:

Field $L$	$\Delta$	$\text{Cl}(\Delta)$
$\mathbb{Q}$	$-3, -4, -7, -8, -11, -12, -16, -19, -27, -28, -43, -67, -163$	trivial
$\mathbb{Q}(\sqrt{2})$	$-24, -32, -64, -88$	$\mathbb{Z}/2\mathbb{Z}$
$\mathbb{Q}(\sqrt{3})$	$-36, -48$	$\mathbb{Z}/2\mathbb{Z}$
$\mathbb{Q}(\sqrt{5})$	$-15, -20, -35, -40, -60, -75, -100, -115, -235$	$\mathbb{Z}/2\mathbb{Z}$
$\mathbb{Q}(\sqrt{13})$	$-52, -91, -403$	$\mathbb{Z}/2\mathbb{Z}$
$\mathbb{Q}(\sqrt{17})$	$-51, -187$	$\mathbb{Z}/2\mathbb{Z}$
$\mathbb{Q}(\sqrt{2}, \sqrt{3})$	$-96, -192, -288$	$(\mathbb{Z}/2\mathbb{Z})^2$
$\mathbb{Q}(\sqrt{3}, \sqrt{5})$	$-180, -240$	$(\mathbb{Z}/2\mathbb{Z})^2$
$\mathbb{Q}(\sqrt{5}, \sqrt{13})$	$-195, -520, -715$	$(\mathbb{Z}/2\mathbb{Z})^2$
$\mathbb{Q}(\sqrt{2}, \sqrt{5})$	$-120, -160, -280, -760$	$(\mathbb{Z}/2\mathbb{Z})^2$
$\mathbb{Q}(\sqrt{5}, \sqrt{17})$	$-340, -595$	$(\mathbb{Z}/2\mathbb{Z})^2$
$\mathbb{Q}(\sqrt{2}, \sqrt{3}, \sqrt{5})$	$-480, -960$	$(\mathbb{Z}/2\mathbb{Z})^3$

- ▶ If  $\mathbb{Q}(\tau_1) = \mathbb{Q}(\tau_2)$  then  $\Delta_1/\Delta_2 \in \{1, 4, 1/4\}$  or  $\Delta_1, \Delta_2 \in \{-3, -12, -27\}$ .



# The Proof

**May assume:**

$$\blacktriangleright \tau_i = \frac{-b_i + \sqrt{\Delta_i}}{2a_i};$$

# The Proof

**May assume:**

- ▶  $\tau_i = \frac{-b_i + \sqrt{\Delta_i}}{2a_i}$ ;
- ▶  $\mathbb{Q}(j(\tau_1)) = \mathbb{Q}(j(\tau_2)) = L$

# The Proof

**May assume:**

- ▶  $\tau_i = \frac{-b_i + \sqrt{\Delta_i}}{2a_i}$ ;
- ▶  $\mathbb{Q}(j(\tau_1)) = \mathbb{Q}(j(\tau_2)) = L$
- ▶  $[L : \mathbb{Q}] = h(\Delta_1) = h(\Delta_2) \geq 3$

# The Proof

**May assume:**

- ▶  $\tau_i = \frac{-b_i + \sqrt{\Delta_i}}{2a_i}$ ;
- ▶  $\mathbb{Q}(j(\tau_1)) = \mathbb{Q}(j(\tau_2)) = L$
- ▶  $[L : \mathbb{Q}] = h(\Delta_1) = h(\Delta_2) \geq 3$
- ▶  $j(\tau_1) \neq j(\tau_2)$ .

# The Proof

**May assume:**

- ▶  $\tau_i = \frac{-b_i + \sqrt{\Delta_i}}{2a_i}$ ;
- ▶  $\mathbb{Q}(j(\tau_1)) = \mathbb{Q}(j(\tau_2)) = L$
- ▶  $[L : \mathbb{Q}] = h(\Delta_1) = h(\Delta_2) \geq 3$
- ▶  $j(\tau_1) \neq j(\tau_2)$ .

# The Proof

## May assume:

- ▶  $\tau_i = \frac{-b_i + \sqrt{\Delta_i}}{2a_i}$ ;
- ▶  $\mathbb{Q}(j(\tau_1)) = \mathbb{Q}(j(\tau_2)) = L$
- ▶  $[L : \mathbb{Q}] = h(\Delta_1) = h(\Delta_2) \geq 3$
- ▶  $j(\tau_1) \neq j(\tau_2)$ .

## Crucial steps:

- ▶  $\mathbb{Q}(\tau_1) = \mathbb{Q}(\tau_2)$ ;

# The Proof

## May assume:

- ▶  $\tau_i = \frac{-b_i + \sqrt{\Delta_i}}{2a_i}$ ;
- ▶  $\mathbb{Q}(j(\tau_1)) = \mathbb{Q}(j(\tau_2)) = L$
- ▶  $[L : \mathbb{Q}] = h(\Delta_1) = h(\Delta_2) \geq 3$
- ▶  $j(\tau_1) \neq j(\tau_2)$ .

## Crucial steps:

- ▶  $\mathbb{Q}(\tau_1) = \mathbb{Q}(\tau_2)$ ;
- ▶  $a_1 = a_2 = 1$ ;

# The Proof

## May assume:

- ▶  $\tau_i = \frac{-b_i + \sqrt{\Delta_i}}{2a_i}$ ;
- ▶  $\mathbb{Q}(j(\tau_1)) = \mathbb{Q}(j(\tau_2)) = L$
- ▶  $[L : \mathbb{Q}] = h(\Delta_1) = h(\Delta_2) \geq 3$
- ▶  $j(\tau_1) \neq j(\tau_2)$ .

## Crucial steps:

- ▶  $\mathbb{Q}(\tau_1) = \mathbb{Q}(\tau_2)$ ;
- ▶  $a_1 = a_2 = 1$ ;



# The Proof

## May assume:

- ▶  $\tau_i = \frac{-b_i + \sqrt{\Delta_i}}{2a_i}$ ;
- ▶  $\mathbb{Q}(j(\tau_1)) = \mathbb{Q}(j(\tau_2)) = L$
- ▶  $[L : \mathbb{Q}] = h(\Delta_1) = h(\Delta_2) \geq 3$
- ▶  $j(\tau_1) \neq j(\tau_2)$ .

## Crucial steps:

- ▶  $\mathbb{Q}(\tau_1) = \mathbb{Q}(\tau_2)$ ;
- ▶  $a_1 = a_2 = 1$ ;

## Consequences:

- ▶  $\Delta_1 = \Delta$ ,  $\Delta_2 = \Delta$  or  $4\Delta$ .

# The Proof

## May assume:

- ▶  $\tau_i = \frac{-b_i + \sqrt{\Delta_i}}{2a_i}$ ;
- ▶  $\mathbb{Q}(j(\tau_1)) = \mathbb{Q}(j(\tau_2)) = L$
- ▶  $[L : \mathbb{Q}] = h(\Delta_1) = h(\Delta_2) \geq 3$
- ▶  $j(\tau_1) \neq j(\tau_2)$ .

## Crucial steps:

- ▶  $\mathbb{Q}(\tau_1) = \mathbb{Q}(\tau_2)$ ;
- ▶  $a_1 = a_2 = 1$ ;

## Consequences:

- ▶  $\Delta_1 = \Delta$ ,  $\Delta_2 = \Delta$  or  $4\Delta$ .
- ▶  $\tau_1 = \frac{-b_1 + \sqrt{\Delta}}{2}$ ,  $\tau_2 = \frac{-b_2 + \sqrt{\Delta}}{2}$  or  $\tau_2 = \frac{-b_2 + 2\sqrt{\Delta}}{2}$

# The Proof

## May assume:

- ▶  $\tau_i = \frac{-b_i + \sqrt{\Delta_i}}{2a_i}$ ;
- ▶  $\mathbb{Q}(j(\tau_1)) = \mathbb{Q}(j(\tau_2)) = L$
- ▶  $[L : \mathbb{Q}] = h(\Delta_1) = h(\Delta_2) \geq 3$
- ▶  $j(\tau_1) \neq j(\tau_2)$ .

## Crucial steps:

- ▶  $\mathbb{Q}(\tau_1) = \mathbb{Q}(\tau_2)$ ;
- ▶  $a_1 = a_2 = 1$ ;

## Consequences:

- ▶  $\Delta_1 = \Delta$ ,  $\Delta_2 = \Delta$  or  $4\Delta$ .
- ▶  $\tau_1 = \frac{-b_1 + \sqrt{\Delta}}{2}$ ,  $\tau_2 = \frac{-b_2 + \sqrt{\Delta}}{2}$  or  $\tau_2 = \frac{-b_2 + 2\sqrt{\Delta}}{2}$
- ▶ In the first case  $j(\tau_1) = j(\tau_2)$

# The Proof

## May assume:

- ▶  $\tau_i = \frac{-b_i + \sqrt{\Delta_i}}{2a_i}$ ;
- ▶  $\mathbb{Q}(j(\tau_1)) = \mathbb{Q}(j(\tau_2)) = L$
- ▶  $[L : \mathbb{Q}] = h(\Delta_1) = h(\Delta_2) \geq 3$
- ▶  $j(\tau_1) \neq j(\tau_2)$ .

## Crucial steps:

- ▶  $\mathbb{Q}(\tau_1) = \mathbb{Q}(\tau_2)$ ;
- ▶  $a_1 = a_2 = 1$ ;

## Consequences:

- ▶  $\Delta_1 = \Delta$ ,  $\Delta_2 = \Delta$  or  $4\Delta$ .
- ▶  $\tau_1 = \frac{-b_1 + \sqrt{\Delta}}{2}$ ,  $\tau_2 = \frac{-b_2 + \sqrt{\Delta}}{2}$  or  $\tau_2 = \frac{-b_2 + 2\sqrt{\Delta}}{2}$
- ▶ In the first case  $j(\tau_1) = j(\tau_2)$
- ▶ In the second case  $(j(\tau_1), j(\tau_2)) \in Y_0(2)$ .

## The Proof (continued)

- ▶  $Y_0(2)$  is curve of degree 4

## The Proof (continued)

- ▶  $Y_0(2)$  is curve of degree 4
- ▶ Hence  $|Y_0(2) \cap (\text{straight line})| \leq 4$

## The Proof (continued)

- ▶  $Y_0(2)$  is curve of degree 4
- ▶ Hence  $|Y_0(2) \cap (\text{straight line})| \leq 4$
- ▶ Hence  $3 \leq h(\Delta) = h(4\Delta) \leq 4$

## The Proof (continued)

- ▶  $Y_0(2)$  is curve of degree 4
- ▶ Hence  $|Y_0(2) \cap (\text{straight line})| \leq 4$
- ▶ Hence  $3 \leq h(\Delta) = h(4\Delta) \leq 4$
- ▶ There is only five  $\Delta$  with this property:



## The Proof (continued)

- ▶  $Y_0(2)$  is curve of degree 4
- ▶ Hence  $|Y_0(2) \cap (\text{straight line})| \leq 4$
- ▶ Hence  $3 \leq h(\Delta) = h(4\Delta) \leq 4$
- ▶ There is only five  $\Delta$  with this property:

## The Proof (continued)

- ▶  $Y_0(2)$  is curve of degree 4
- ▶ Hence  $|Y_0(2) \cap (\text{straight line})| \leq 4$
- ▶ Hence  $3 \leq h(\Delta) = h(4\Delta) \leq 4$
- ▶ There is only five  $\Delta$  with this property:

$h = 3$ :  $-23, -31$ ;

## The Proof (continued)

- ▶  $Y_0(2)$  is curve of degree 4
- ▶ Hence  $|Y_0(2) \cap (\text{straight line})| \leq 4$
- ▶ Hence  $3 \leq h(\Delta) = h(4\Delta) \leq 4$
- ▶ There is only five  $\Delta$  with this property:

$h = 3$ :  $-23, -31$ ;

$h = 4$ :  $-7 \cdot 3^2, -39, -55$ .

## The Proof (continued)

- ▶  $Y_0(2)$  is curve of degree 4
- ▶ Hence  $|Y_0(2) \cap (\text{straight line})| \leq 4$
- ▶ Hence  $3 \leq h(\Delta) = h(4\Delta) \leq 4$
- ▶ There is only five  $\Delta$  with this property:

$h = 3$ :  $-23, -31$ ;

$h = 4$ :  $-7 \cdot 3^2, -39, -55$ .

## The Proof (continued)

- ▶  $Y_0(2)$  is curve of degree 4
- ▶ Hence  $|Y_0(2) \cap (\text{straight line})| \leq 4$
- ▶ Hence  $3 \leq h(\Delta) = h(4\Delta) \leq 4$
- ▶ There is only five  $\Delta$  with this property:

$$h = 3: -23, -31;$$

$$h = 4: -7 \cdot 3^2, -39, -55.$$

One rules them out using PARI.



## Complex Multiplication

Lattices

$j$ -invariant

Complex Multiplication

Class Field Theory

The Class Number

## Theorem of André

Special Points and Special Curves

Theorem of André

## CM-Points on Straight Lines

Kühne's "uniformity observation"

CM-Points on Straight Lines

## The Proof

Equality of CM-fields

The Proof

## Proof of Theorem ECMF

Discriminants with Class Group Annihilated by 2

Proof of Theorem ECMF

# Discriminants with Class Group Annihilated by 2

## Known $\Delta$ with $\text{Cl}(\Delta)^2 = 1$

- 3,  $-3 \cdot 2^2$ ,  $-3 \cdot 3^2$ ,  $-3 \cdot 4^2$ ,  $-3 \cdot 5^2$ ,  $-3 \cdot 7^2$ ,  $-3 \cdot 8^2$ , -4,  $-4 \cdot 2^2$ ,  $-4 \cdot 3^2$ ,  $-4 \cdot 4^2$ ,  $-4 \cdot 5^2$ ,  
- 7,  $-7 \cdot 2^2$ ,  $-7 \cdot 4^2$ ,  $-7 \cdot 8^2$ , -8,  $-8 \cdot 2^2$ ,  $-8 \cdot 3^2$ ,  $-8 \cdot 6^2$ , -11,  $-11 \cdot 3^2$ ,  
- 15,  $-15 \cdot 2^2$ ,  $-15 \cdot 4^2$ ,  $-15 \cdot 8^2$ , -19, -20,  $-20 \cdot 3^2$ , -24,  $-24 \cdot 2^2$ , -35,  $-35 \cdot 3^2$ , -40,  $-40 \cdot 2^2$ ,  
- 43, -51, -52, -67, -84, -88,  $-88 \cdot 2^2$ , -91, -115, -120,  $-120 \cdot 2^2$ , -123, -132, -148, -163,  
- 168,  $-168 \cdot 2^2$ , -187, -195, -228, -232,  $-232 \cdot 2^2$ , -235, -267, -280,  $-280 \cdot 2^2$ , -312,  $-312 \cdot 2^2$ ,  
- 340, -372, -403, -408,  $-408 \cdot 2^2$ , -420, -427, -435, -483, -520,  $-520 \cdot 2^2$ , -532, -555, -595,  
- 627, -660, -708, -715, -760,  $-760 \cdot 2^2$ , -795, -840,  $-840 \cdot 2^2$ , -1012, -1092, -1155,  
- 1320,  $-1320 \cdot 2^2$ , -1380, -1428, -1435, -1540, -1848,  $-1848 \cdot 2^2$ , -1995, -3003, -3315, -5460.

# Discriminants with Class Group Annihilated by 2

## Known $\Delta$ with $\text{Cl}(\Delta)^2 = 1$

$-3, -3 \cdot 2^2, -3 \cdot 3^2, -3 \cdot 4^2, -3 \cdot 5^2, -3 \cdot 7^2, -3 \cdot 8^2, -4, -4 \cdot 2^2, -4 \cdot 3^2, -4 \cdot 4^2, -4 \cdot 5^2,$   
 $-7, -7 \cdot 2^2, -7 \cdot 4^2, -7 \cdot 8^2, -8, -8 \cdot 2^2, -8 \cdot 3^2, -8 \cdot 6^2, -11, -11 \cdot 3^2,$   
 $-15, -15 \cdot 2^2, -15 \cdot 4^2, -15 \cdot 8^2, -19, -20, -20 \cdot 3^2, -24, -24 \cdot 2^2, -35, -35 \cdot 3^2, -40, -40 \cdot 2^2,$   
 $-43, -51, -52, -67, -84, -88, -88 \cdot 2^2, -91, -115, -120, -120 \cdot 2^2, -123, -132, -148, -163,$   
 $-168, -168 \cdot 2^2, -187, -195, -228, -232, -232 \cdot 2^2, -235, -267, -280, -280 \cdot 2^2, -312, -312 \cdot 2^2,$   
 $-340, -372, -403, -408, -408 \cdot 2^2, -420, -427, -435, -483, -520, -520 \cdot 2^2, -532, -555, -595,$   
 $-627, -660, -708, -715, -760, -760 \cdot 2^2, -795, -840, -840 \cdot 2^2, -1012, -1092, -1155,$   
 $-1320, -1320 \cdot 2^2, -1380, -1428, -1435, -1540, -1848, -1848 \cdot 2^2, -1995, -3003, -3315, -5460.$

► **Weinberger** (1973): All field discriminants  $D$  with  $\text{Cl}(D)^2 = 1$  belong to the list above



# Discriminants with Class Group Annihilated by 2

## Known $\Delta$ with $\text{Cl}(\Delta)^2 = 1$

$-3, -3 \cdot 2^2, -3 \cdot 3^2, -3 \cdot 4^2, -3 \cdot 5^2, -3 \cdot 7^2, -3 \cdot 8^2, -4, -4 \cdot 2^2, -4 \cdot 3^2, -4 \cdot 4^2, -4 \cdot 5^2,$   
 $-7, -7 \cdot 2^2, -7 \cdot 4^2, -7 \cdot 8^2, -8, -8 \cdot 2^2, -8 \cdot 3^2, -8 \cdot 6^2, -11, -11 \cdot 3^2,$   
 $-15, -15 \cdot 2^2, -15 \cdot 4^2, -15 \cdot 8^2, -19, -20, -20 \cdot 3^2, -24, -24 \cdot 2^2, -35, -35 \cdot 3^2, -40, -40 \cdot 2^2,$   
 $-43, -51, -52, -67, -84, -88, -88 \cdot 2^2, -91, -115, -120, -120 \cdot 2^2, -123, -132, -148, -163,$   
 $-168, -168 \cdot 2^2, -187, -195, -228, -232, -232 \cdot 2^2, -235, -267, -280, -280 \cdot 2^2, -312, -312 \cdot 2^2,$   
 $-340, -372, -403, -408, -408 \cdot 2^2, -420, -427, -435, -483, -520, -520 \cdot 2^2, -532, -555, -595,$   
 $-627, -660, -708, -715, -760, -760 \cdot 2^2, -795, -840, -840 \cdot 2^2, -1012, -1092, -1155,$   
 $-1320, -1320 \cdot 2^2, -1380, -1428, -1435, -1540, -1848, -1848 \cdot 2^2, -1995, -3003, -3315, -5460.$

► **Weinberger** (1973): All field discriminants  $D$  with  $\text{Cl}(D)^2 = 1$  belong to the list above

# Discriminants with Class Group Annihilated by 2

## Known $\Delta$ with $\text{Cl}(\Delta)^2 = 1$

$-3, -3 \cdot 2^2, -3 \cdot 3^2, -3 \cdot 4^2, -3 \cdot 5^2, -3 \cdot 7^2, -3 \cdot 8^2, -4, -4 \cdot 2^2, -4 \cdot 3^2, -4 \cdot 4^2, -4 \cdot 5^2,$   
 $-7, -7 \cdot 2^2, -7 \cdot 4^2, -7 \cdot 8^2, -8, -8 \cdot 2^2, -8 \cdot 3^2, -8 \cdot 6^2, -11, -11 \cdot 3^2,$   
 $-15, -15 \cdot 2^2, -15 \cdot 4^2, -15 \cdot 8^2, -19, -20, -20 \cdot 3^2, -24, -24 \cdot 2^2, -35, -35 \cdot 3^2, -40, -40 \cdot 2^2,$   
 $-43, -51, -52, -67, -84, -88, -88 \cdot 2^2, -91, -115, -120, -120 \cdot 2^2, -123, -132, -148, -163,$   
 $-168, -168 \cdot 2^2, -187, -195, -228, -232, -232 \cdot 2^2, -235, -267, -280, -280 \cdot 2^2, -312, -312 \cdot 2^2,$   
 $-340, -372, -403, -408, -408 \cdot 2^2, -420, -427, -435, -483, -520, -520 \cdot 2^2, -532, -555, -595,$   
 $-627, -660, -708, -715, -760, -760 \cdot 2^2, -795, -840, -840 \cdot 2^2, -1012, -1092, -1155,$   
 $-1320, -1320 \cdot 2^2, -1380, -1428, -1435, -1540, -1848, -1848 \cdot 2^2, -1995, -3003, -3315, -5460.$

- ▶ **Weinberger** (1973): All field discriminants  $D$  with  $\text{Cl}(D)^2 = 1$  belong to the list above **with at most one exception**.

# Discriminants with Class Group Annihilated by 2

## Known $\Delta$ with $\text{Cl}(\Delta)^2 = 1$

- 3,  $-3 \cdot 2^2$ ,  $-3 \cdot 3^2$ ,  $-3 \cdot 4^2$ ,  $-3 \cdot 5^2$ ,  $-3 \cdot 7^2$ ,  $-3 \cdot 8^2$ , -4,  $-4 \cdot 2^2$ ,  $-4 \cdot 3^2$ ,  $-4 \cdot 4^2$ ,  $-4 \cdot 5^2$ ,  
- 7,  $-7 \cdot 2^2$ ,  $-7 \cdot 4^2$ ,  $-7 \cdot 8^2$ , -8,  $-8 \cdot 2^2$ ,  $-8 \cdot 3^2$ ,  $-8 \cdot 6^2$ , -11,  $-11 \cdot 3^2$ ,  
- 15,  $-15 \cdot 2^2$ ,  $-15 \cdot 4^2$ ,  $-15 \cdot 8^2$ , -19, -20,  $-20 \cdot 3^2$ , -24,  $-24 \cdot 2^2$ , -35,  $-35 \cdot 3^2$ , -40,  $-40 \cdot 2^2$ ,  
- 43, -51, -52, -67, -84, -88,  $-88 \cdot 2^2$ , -91, -115, -120,  $-120 \cdot 2^2$ , -123, -132, -148, -163,  
- 168,  $-168 \cdot 2^2$ , -187, -195, -228, -232,  $-232 \cdot 2^2$ , -235, -267, -280,  $-280 \cdot 2^2$ , -312,  $-312 \cdot 2^2$ ,  
- 340, -372, -403, -408,  $-408 \cdot 2^2$ , -420, -427, -435, -483, -520,  $-520 \cdot 2^2$ , -532, -555, -595,  
- 627, -660, -708, -715, -760,  $-760 \cdot 2^2$ , -795, -840,  $-840 \cdot 2^2$ , -1012, -1092, -1155,  
- 1320,  $-1320 \cdot 2^2$ , -1380, -1428, -1435, -1540, -1848,  $-1848 \cdot 2^2$ , -1995, -3003, -3315, -5460.

- ▶ **Weinberger** (1973): All field discriminants  $D$  with  $\text{Cl}(D)^2 = 1$  belong to the list above **with at most one exception**.
- ▶ **Corollary:** There exists  $D^*$  such that: if  $\Delta = Df^2$  with  $\text{Cl}(\Delta)^2 = 1$  is **not** in the list then  $D = D^*$ .

# Discriminants with Class Group Annihilated by 2

## Known $\Delta$ with $\text{Cl}(\Delta)^2 = 1$

- 3,  $-3 \cdot 2^2$ ,  $-3 \cdot 3^2$ ,  $-3 \cdot 4^2$ ,  $-3 \cdot 5^2$ ,  $-3 \cdot 7^2$ ,  $-3 \cdot 8^2$ , -4,  $-4 \cdot 2^2$ ,  $-4 \cdot 3^2$ ,  $-4 \cdot 4^2$ ,  $-4 \cdot 5^2$ ,  
- 7,  $-7 \cdot 2^2$ ,  $-7 \cdot 4^2$ ,  $-7 \cdot 8^2$ , -8,  $-8 \cdot 2^2$ ,  $-8 \cdot 3^2$ ,  $-8 \cdot 6^2$ , -11,  $-11 \cdot 3^2$ ,  
- 15,  $-15 \cdot 2^2$ ,  $-15 \cdot 4^2$ ,  $-15 \cdot 8^2$ , -19, -20,  $-20 \cdot 3^2$ , -24,  $-24 \cdot 2^2$ , -35,  $-35 \cdot 3^2$ , -40,  $-40 \cdot 2^2$ ,  
- 43, -51, -52, -67, -84, -88,  $-88 \cdot 2^2$ , -91, -115, -120,  $-120 \cdot 2^2$ , -123, -132, -148, -163,  
- 168,  $-168 \cdot 2^2$ , -187, -195, -228, -232,  $-232 \cdot 2^2$ , -235, -267, -280,  $-280 \cdot 2^2$ , -312,  $-312 \cdot 2^2$ ,  
- 340, -372, -403, -408,  $-408 \cdot 2^2$ , -420, -427, -435, -483, -520,  $-520 \cdot 2^2$ , -532, -555, -595,  
- 627, -660, -708, -715, -760,  $-760 \cdot 2^2$ , -795, -840,  $-840 \cdot 2^2$ , -1012, -1092, -1155,  
- 1320,  $-1320 \cdot 2^2$ , -1380, -1428, -1435, -1540, -1848,  $-1848 \cdot 2^2$ , -1995, -3003, -3315, -5460.

- ▶ **Weinberger** (1973): All field discriminants  $D$  with  $\text{Cl}(D)^2 = 1$  belong to the list above **with at most one exception**.
- ▶ **Corollary:** There exists  $D^*$  such that: if  $\Delta = Df^2$  with  $\text{Cl}(\Delta)^2 = 1$  is **not** in the list then  $D = D^*$ .
- ▶ Class numbers of discriminants from the list are at most 16.

# Discriminants with Class Group Annihilated by 2

## Known $\Delta$ with $\text{Cl}(\Delta)^2 = 1$

- 3,  $-3 \cdot 2^2$ ,  $-3 \cdot 3^2$ ,  $-3 \cdot 4^2$ ,  $-3 \cdot 5^2$ ,  $-3 \cdot 7^2$ ,  $-3 \cdot 8^2$ , -4,  $-4 \cdot 2^2$ ,  $-4 \cdot 3^2$ ,  $-4 \cdot 4^2$ ,  $-4 \cdot 5^2$ ,  
- 7,  $-7 \cdot 2^2$ ,  $-7 \cdot 4^2$ ,  $-7 \cdot 8^2$ , -8,  $-8 \cdot 2^2$ ,  $-8 \cdot 3^2$ ,  $-8 \cdot 6^2$ , -11,  $-11 \cdot 3^2$ ,  
- 15,  $-15 \cdot 2^2$ ,  $-15 \cdot 4^2$ ,  $-15 \cdot 8^2$ , -19, -20,  $-20 \cdot 3^2$ , -24,  $-24 \cdot 2^2$ , -35,  $-35 \cdot 3^2$ , -40,  $-40 \cdot 2^2$ ,  
- 43, -51, -52, -67, -84, -88,  $-88 \cdot 2^2$ , -91, -115, -120,  $-120 \cdot 2^2$ , -123, -132, -148, -163,  
- 168,  $-168 \cdot 2^2$ , -187, -195, -228, -232,  $-232 \cdot 2^2$ , -235, -267, -280,  $-280 \cdot 2^2$ , -312,  $-312 \cdot 2^2$ ,  
- 340, -372, -403, -408,  $-408 \cdot 2^2$ , -420, -427, -435, -483, -520,  $-520 \cdot 2^2$ , -532, -555, -595,  
- 627, -660, -708, -715, -760,  $-760 \cdot 2^2$ , -795, -840,  $-840 \cdot 2^2$ , -1012, -1092, -1155,  
- 1320,  $-1320 \cdot 2^2$ , -1380, -1428, -1435, -1540, -1848,  $-1848 \cdot 2^2$ , -1995, -3003, -3315, -5460.

- ▶ **Weinberger** (1973): All field discriminants  $D$  with  $\text{Cl}(D)^2 = 1$  belong to the list above **with at most one exception**.
- ▶ **Corollary:** There exists  $D^*$  such that: if  $\Delta = Df^2$  with  $\text{Cl}(\Delta)^2 = 1$  is **not** in the list then  $D = D^*$ .
- ▶ Class numbers of discriminants from the list are at most 16.
- ▶ **Watkins** (2006): the list contains all  $\Delta$  with  $|\text{Cl}(\Delta)^2| = 1$  and  $h(\Delta) \leq 64$ .

# Discriminants with Class Group Annihilated by 2

## Known $\Delta$ with $\text{Cl}(\Delta)^2 = 1$

- 3,  $-3 \cdot 2^2$ ,  $-3 \cdot 3^2$ ,  $-3 \cdot 4^2$ ,  $-3 \cdot 5^2$ ,  $-3 \cdot 7^2$ ,  $-3 \cdot 8^2$ , -4,  $-4 \cdot 2^2$ ,  $-4 \cdot 3^2$ ,  $-4 \cdot 4^2$ ,  $-4 \cdot 5^2$ ,  
- 7,  $-7 \cdot 2^2$ ,  $-7 \cdot 4^2$ ,  $-7 \cdot 8^2$ , -8,  $-8 \cdot 2^2$ ,  $-8 \cdot 3^2$ ,  $-8 \cdot 6^2$ , -11,  $-11 \cdot 3^2$ ,  
- 15,  $-15 \cdot 2^2$ ,  $-15 \cdot 4^2$ ,  $-15 \cdot 8^2$ , -19, -20,  $-20 \cdot 3^2$ , -24,  $-24 \cdot 2^2$ , -35,  $-35 \cdot 3^2$ , -40,  $-40 \cdot 2^2$ ,  
- 43, -51, -52, -67, -84, -88,  $-88 \cdot 2^2$ , -91, -115, -120,  $-120 \cdot 2^2$ , -123, -132, -148, -163,  
- 168,  $-168 \cdot 2^2$ , -187, -195, -228, -232,  $-232 \cdot 2^2$ , -235, -267, -280,  $-280 \cdot 2^2$ , -312,  $-312 \cdot 2^2$ ,  
- 340, -372, -403, -408,  $-408 \cdot 2^2$ , -420, -427, -435, -483, -520,  $-520 \cdot 2^2$ , -532, -555, -595,  
- 627, -660, -708, -715, -760,  $-760 \cdot 2^2$ , -795, -840,  $-840 \cdot 2^2$ , -1012, -1092, -1155,  
- 1320,  $-1320 \cdot 2^2$ , -1380, -1428, -1435, -1540, -1848,  $-1848 \cdot 2^2$ , -1995, -3003, -3315, -5460.

- ▶ **Weinberger** (1973): All field discriminants  $D$  with  $\text{Cl}(D)^2 = 1$  belong to the list above **with at most one exception**.
- ▶ **Corollary:** There exists  $D^*$  such that: if  $\Delta = Df^2$  with  $\text{Cl}(\Delta)^2 = 1$  is **not** in the list then  $D = D^*$ .
- ▶ Class numbers of discriminants from the list are at most 16.
- ▶ **Watkins** (2006): the list contains all  $\Delta$  with  $|\text{Cl}(\Delta)^2| = 1$  and  $h(\Delta) \leq 64$ .
- ▶ Hence: if  $\Delta$  with  $\text{Cl}(\Delta)^2 = 1$  is **not** in the list then  $h(\Delta) \geq 128$ .

# Proof of Theorem ECMF

- ▶ Assume that  $Q(\tau_1) \neq Q(\tau_2)$  and  $Q(j(\tau_1)) = Q(j(\tau_2))$ .


# Proof of Theorem ECMF

- ▶ Assume that  $\mathbb{Q}(\tau_1) \neq \mathbb{Q}(\tau_2)$  and  $\mathbb{Q}(j(\tau_1)) = \mathbb{Q}(j(\tau_2))$ .
- ▶ Set  $M = \mathbb{Q}(\tau_1, \tau_2, j(\tau_1)) = \mathbb{Q}(\tau_1, \tau_2, j(\tau_2))$ .




# Proof of Theorem ECMF

- ▶ Assume that  $\mathbb{Q}(\tau_1) \neq \mathbb{Q}(\tau_2)$  and  $\mathbb{Q}(j(\tau_1)) = \mathbb{Q}(j(\tau_2))$ .
- ▶ Set  $M = \mathbb{Q}(\tau_1, \tau_2, j(\tau_1)) = \mathbb{Q}(\tau_1, \tau_2, j(\tau_2))$ .
- ▶ **André, Edixhoven** (1998):  $G = \text{Gal}(M/\mathbb{Q}(\tau_1, \tau_2))$  is annihilated by 2.



# Proof of Theorem ECMF


- ▶ Assume that  $\mathbb{Q}(\tau_1) \neq \mathbb{Q}(\tau_2)$  and  $\mathbb{Q}(j(\tau_1)) = \mathbb{Q}(j(\tau_2))$ .
- ▶ Set  $M = \mathbb{Q}(\tau_1, \tau_2, j(\tau_1)) = \mathbb{Q}(\tau_1, \tau_2, j(\tau_2))$ .
- ▶ **André, Edixhoven** (1998):  $G = \text{Gal}(M/\mathbb{Q}(\tau_1, \tau_2))$  is annihilated by 2.



- ▶ Consequence: each  $\text{Cl}(\Delta_i)$  is of type  $\mathbb{Z}/4 \times \mathbb{Z}/2 \times \cdots \times \mathbb{Z}/2$  or  $\mathbb{Z}/2 \times \cdots \times \mathbb{Z}/2$ .

# Proof of Theorem ECMF


- ▶ Assume that  $\mathbb{Q}(\tau_1) \neq \mathbb{Q}(\tau_2)$  and  $\mathbb{Q}(j(\tau_1)) = \mathbb{Q}(j(\tau_2))$ .
- ▶ Set  $M = \mathbb{Q}(\tau_1, \tau_2, j(\tau_1)) = \mathbb{Q}(\tau_1, \tau_2, j(\tau_2))$ .
- ▶ **André, Edixhoven** (1998):  $G = \text{Gal}(M/\mathbb{Q}(\tau_1, \tau_2))$  is annihilated by 2.



- ▶ Consequence: each  $\text{Cl}(\Delta_i)$  is of type  $\mathbb{Z}/4 \times \mathbb{Z}/2 \times \dots \times \mathbb{Z}/2$  or  $\mathbb{Z}/2 \times \dots \times \mathbb{Z}/2$ .
- ▶ Simple group theory: first type is impossible. Hence both  $\text{Cl}(\Delta_i)$  are annihilated by 2.

# Proof of Theorem ECMF


- ▶ Assume that  $\mathbb{Q}(\tau_1) \neq \mathbb{Q}(\tau_2)$  and  $\mathbb{Q}(j(\tau_1)) = \mathbb{Q}(j(\tau_2))$ .
- ▶ Set  $M = \mathbb{Q}(\tau_1, \tau_2, j(\tau_1)) = \mathbb{Q}(\tau_1, \tau_2, j(\tau_2))$ .
- ▶ **André, Edixhoven** (1998):  $G = \text{Gal}(M/\mathbb{Q}(\tau_1, \tau_2))$  is annihilated by 2.



- ▶ Consequence: each  $\text{Cl}(\Delta_i)$  is of type  $\mathbb{Z}/4 \times \mathbb{Z}/2 \times \dots \times \mathbb{Z}/2$  or  $\mathbb{Z}/2 \times \dots \times \mathbb{Z}/2$ .
- ▶ Simple group theory: first type is impossible. Hence both  $\text{Cl}(\Delta_i)$  are annihilated by 2.
- ▶ Weinberger: since  $D_1 \neq D_2$

# Proof of Theorem ECMF


- ▶ Assume that  $\mathbb{Q}(\tau_1) \neq \mathbb{Q}(\tau_2)$  and  $\mathbb{Q}(j(\tau_1)) = \mathbb{Q}(j(\tau_2))$ .
- ▶ Set  $M = \mathbb{Q}(\tau_1, \tau_2, j(\tau_1)) = \mathbb{Q}(\tau_1, \tau_2, j(\tau_2))$ .
- ▶ **André, Edixhoven** (1998):  $G = \text{Gal}(M/\mathbb{Q}(\tau_1, \tau_2))$  is annihilated by 2.



- ▶ Consequence: each  $\text{Cl}(\Delta_i)$  is of type  $\mathbb{Z}/4 \times \mathbb{Z}/2 \times \dots \times \mathbb{Z}/2$  or  $\mathbb{Z}/2 \times \dots \times \mathbb{Z}/2$ .
- ▶ Simple group theory: first type is impossible. Hence both  $\text{Cl}(\Delta_i)$  are annihilated by 2.
- ▶ Weinberger: since  $D_1 \neq D_2$

# Proof of Theorem ECMF


- ▶ Assume that  $\mathbb{Q}(\tau_1) \neq \mathbb{Q}(\tau_2)$  and  $\mathbb{Q}(j(\tau_1)) = \mathbb{Q}(j(\tau_2))$ .
- ▶ Set  $M = \mathbb{Q}(\tau_1, \tau_2, j(\tau_1)) = \mathbb{Q}(\tau_1, \tau_2, j(\tau_2))$ .
- ▶ **André, Edixhoven** (1998):  $G = \text{Gal}(M/\mathbb{Q}(\tau_1, \tau_2))$  is annihilated by 2.



- ▶ Consequence: each  $\text{Cl}(\Delta_i)$  is of type  $\mathbb{Z}/4 \times \mathbb{Z}/2 \times \dots \times \mathbb{Z}/2$  or  $\mathbb{Z}/2 \times \dots \times \mathbb{Z}/2$ .
- ▶ Simple group theory: first type is impossible. Hence both  $\text{Cl}(\Delta_i)$  are annihilated by 2.
- ▶ Weinberger: since  $D_1 \neq D_2$  **one of  $\Delta_1, \Delta_2$  is in the list.**

# Proof of Theorem ECMF


- ▶ Assume that  $\mathbb{Q}(\tau_1) \neq \mathbb{Q}(\tau_2)$  and  $\mathbb{Q}(j(\tau_1)) = \mathbb{Q}(j(\tau_2))$ .
- ▶ Set  $M = \mathbb{Q}(\tau_1, \tau_2, j(\tau_1)) = \mathbb{Q}(\tau_1, \tau_2, j(\tau_2))$ .
- ▶ **André, Edixhoven** (1998):  $G = \text{Gal}(M/\mathbb{Q}(\tau_1, \tau_2))$  is annihilated by 2.



- ▶ Consequence: each  $\text{Cl}(\Delta_i)$  is of type  $\mathbb{Z}/4 \times \mathbb{Z}/2 \times \dots \times \mathbb{Z}/2$  or  $\mathbb{Z}/2 \times \dots \times \mathbb{Z}/2$ .
- ▶ Simple group theory: first type is impossible. Hence both  $\text{Cl}(\Delta_i)$  are annihilated by 2.
- ▶ Weinberger: since  $D_1 \neq D_2$  **one of  $\Delta_1, \Delta_2$  is in the list.**
- ▶ Hence the other is in the list as well.

# Proof of Theorem ECMF

- ▶ Assume that  $\mathbb{Q}(\tau_1) \neq \mathbb{Q}(\tau_2)$  and  $\mathbb{Q}(j(\tau_1)) = \mathbb{Q}(j(\tau_2))$ .
- ▶ Set  $M = \mathbb{Q}(\tau_1, \tau_2, j(\tau_1)) = \mathbb{Q}(\tau_1, \tau_2, j(\tau_2))$ .
- ▶ **André, Edixhoven** (1998):  $G = \text{Gal}(M/\mathbb{Q}(\tau_1, \tau_2))$  is annihilated by 2.



- ▶ Consequence: each  $\text{Cl}(\Delta_i)$  is of type  $\mathbb{Z}/4 \times \mathbb{Z}/2 \times \dots \times \mathbb{Z}/2$  or  $\mathbb{Z}/2 \times \dots \times \mathbb{Z}/2$ .
- ▶ Simple group theory: first type is impossible. Hence both  $\text{Cl}(\Delta_i)$  are annihilated by 2.
- ▶ Weinberger: since  $D_1 \neq D_2$  **one of  $\Delta_1, \Delta_2$  is in the list.**
- ▶ Hence the other is in the list as well.
- ▶ Verification with PARI completes the proof.